

STANFORD UNIV

CONTRIBUTIONS
FROM THE
CUSHMAN LABORATORY
FOR
FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 1-5

PRICE \$1.00 POSTPAID

SHARON, MASSACHUSETTS, U. S. A.

CONTRIBUTIONS FROM THE CUSHMAN
LABORATORY FOR FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 1—5 INCLUSIVE

Owing to the demand of those who wish to bind these Contributions, the following Index has been prepared. Volume 3, part 1 already has a generic index, and so only the new genera of that part have been included in the present Index.

All new genera and species appear in bold-faced type, others in italic type. An abbreviated form gives the geologic formation and general locality with each species.

Papers are grouped both by geographic and geologic headings as well as by subject.

In order to conserve space, the following abbreviated form has been used. Volume number with the part number following in parenthesis followed by a colon. After the colon comes the page number, then the plate number in bold-faced type, and the figure number in italics. It is hoped that this will give ready reference to the subject matter of these volumes.

Agathammina, 3(4) : 188; **protea** (Permian-Texas), 4(2) : 43, 5, 6. [Alabama]. Additional Foraminifera from the Eocene of Alabama, 4(3) : 73; Eocene Foraminifera from the Cocoa Sand of Alabama, 1(3) : 65.

Allomorphinella, 3(1) : 86.

Ammobaculites **compressa** (Penn.-Michigan), 3(2) : 108, 22, 2; **inconspicua** (Penn.-Texas), 4(2) : 41, 5, 2; **spirans** (Penn.-Texas), 3(3) : 149, 26, 10; **stenomeca** (Penn.-Texas), 4(2) : 39, 5, 1; **stormi** (Penn.-Texas), 4(2) : 41, 5, 3, 4; **taylorensis** (Cret.-Texas), 5(3) : 64, 10, 6.

Ammochilostoma (?) **triloculina** (Penn.-Texas), 3(3) : 152, 27, 5.

Ammodiscoides **conica** (Penn.-Texas), 4(2) : 39, 4, 10.

Ammodiscus annularis (Penn.-Michigan), 3(2) : 108, 22, 3; **glabratus** (Cret.-Trinidad), 4(4) : 87, 12, 6; **pennyi** (Cret.-Trinidad), 4(4) : 87, 12, 4, 5.
Ammolagena clavata (Cret.-Trinidad), 4(4) : 90, 12, 14.
Ammovertella, 4(1) : 8.
Angulogerina, 3(1) : 69.
Anomalina cocoaensis (Eoc.-Ala.), 4(3) : 75, 10, 4; **velascoensis** (Cret.-Mexico), 1(1) : 21, 3, 3.
Anomalinella, 3(1) : 93.
Apterrinella, 4(3) : 64.
Arenobulimina, 2(4) : 80.
Assilina, 3(4) : 189.
Asterigerina, 3(4) : 190; **bracteata** (Olig.-Miss), 5(2) : 48, 8, 6.
Baggina, 2(3), 63; **californica** (Mioc.-Calif.), 2(3) : 64, 9, 8.
Bathysiphon sp(?) (Tert.-Venezuela), 5(4) : 78, 12, 1.
Bifarina dalli (Eoc.-Ala.), 2(2) : 31, 4, 5; **vicksburgensis** (Olig.-Miss.), 5(2) : 44, 8, 3, 4.
Bigenerina ciscoensis (Penn.-Texas), 4(3) : 63, 8, 5.
Bolivina advena (Mioc.-Calif.), 1(2) : 29, 5, 1; 2(3) : 54; var. **ornata** (Mioc.-Calif.), 1(2) : 29, 5, 2; 2(3) : 54; var. **striatella** (Mioc.-Calif.), 1(2) : 30, 5, 3; 2(3) : 54; **alazanensis** (Alazan-Mexico), 1(4) : 82, 12, 1; **argentea** (Plioc.-Calif.), 2(2) : 42, 6, 5; **attenuata** (Eoc.-Ala.), 2(2) : 30, 4, 4; **brevior** (Mioc.-Calif.), 1(2) : 31, 5, 8; 2(3) : 54; **caelata** (Tert.-Venezuela), 5(4) : 93, 13, 28; **californica** (Mioc.-Calif.), 1(2) : 32, 5, 10; 2(3) : 54; **clavata** (Cret.-Texas), 2(4) : 87, 12, 5; **conica** (Mioc.-Calif.), 1(2) : 30, 5, 4; 2(3) : 54; **costata** (Plioc.-Calif.), 2(2) : 41, 6, 2; var. **bicostata** (Plioc.-Calif.), 2(2) : 42; var. **interjuncta** (Plioc.-Calif.), 2(2) : 41, 6, 3; **decurrrens** (Cret.-Texas), 2(4) : 88, 12, 4; **decurtata** (Plioc.-Calif.), 2(2) : 44, 6, 7; **decussata** (Mioc.-Calif.), 1(2) : 31, 5, 6; 2(3) : 54; **floridana** (Tert.-Venezuela), 5(4) : 93; **gardnerae** (Eoc.-Georgia), 2(2) : 31, 4, 7; **gemma** (Cret.-Texas), 2(4) : 87, 12, 3; **hughesi** (Plioc.-Calif.), 2(2) : 43, 6, 4; **imbricata** (Mioc.-Calif.), 1(2) : 31, 5, 7; 2(3) : 54; **incrassata** (Cret.-Mexico), 2(1) : 19, 2, 1; 2(4) : 86, 12, 1; var. **limonensis** (Cret.-Mexico), 2(1) : 19, 2, 2; 2(4) : 86; **mantaensis** (Tert.-Venezuela), 5(4) : 92, 13, 27; **marginata** (Mioc.-Calif.), 1(2) : 30, 5, 5; 2(3) : 54; **mexicana** (Alazan-Mexico), 1(4) : 81, 12, 2; var. **aliformis** (Alazan-Mexico), 1(4) : 82, 12, 3, 4; var. **horizontalis** (Alazan-Mexico), 1(4) : 82, 12, 5; **pisciformis** (Tert.-Venezuela), 5(4) : 93, 13, 26; **plicata** (Plioc.-Calif.), 2(2) : 40, 6, 1; **pseudobeyrichi** (Plioc.-Calif.), 2(2) : 45; **punctata** (Plioc.-Calif.), 2(2) : 43; **rhomboidea** (Cret.-Mexico), 2(1) :

19, 2, 3; *seminuda* (Plioc.-Calif.), 2(2): 43; *spiralis* (Eoc.-So. Carolina), 2(2): 31, 4, 6; *subadvena* (Plioc.-Calif.), 2(2): 44, 6, 6; var. *spissa* (Plioc.-Calif.), 2(2): 45, 6, 8; *tectiformis* (Alazan-Mexico), 1(4): 83, 12, 6; *tegulata* (Cret.), 2(4): 86, 12, 2; *tongi* (Tert.-Venezuela), 5(4): 93, 13, 29; *tumida* (Mioc.-Calif.), 1(2): 32, 5, 9; 2(3): 54; *velascoensis* (Cret.-Mexico), 2(4): 88; *watersi* (Cret.-Texas), 2(4): 88, 12, 6.

Bolivinnella and Its Species, The Genus, 5(2): 28.

Bolivinnella, 2(4): 79; 5(2): 28; *australis* (Tert.-Australia), 5(2): 32, 5, 6, 7; *folia* (Rec.-Pac.), 5(2): 29, 5, 1, 2; var. *ornata* (Rec.-Australia), 5(2): 32, 5, 3, 4; *margaritacea* (Mioc.-France), 5(2): 33, 5, 5; *subpectinata* (Olig.-Miss.), 5(2): 34, 5, 8; *virgata* (Mioc.-France), 5(2): 33, 5, 9.

Bolivinita, 2(4): 90; *eleyi* (Cret.), 2(4): 91, 12, 11; *planata* (Cret.-Texas), 3(2): 115, 23, 9.

Bolivinoidea, 2(4): 89; *decorata* (Cret.), 2(4): 89, 12, 9; var. *delicatula* (Cret.), 2(4): 90, 12, 8; 4(4): 99, 14, 9; *rhomboidea* (Cret.-Mexico), 2(4): 90, 12, 10; *trinitatensis* (Cret.-Trinidad), 4(4): 99, 14, 10.

Bradyina, 3(4): 189; *millsapensis* (Penn.-Texas), 4(2): 53, 7, 3, 4.

Bulimina inflata (Tert.-Venezuela), 5(4): 94, 13, 31; *jacksonensis* (Eoc.-Mexico), 1(1): 6, 1, 6, 7; 1(3): 65; var. *cuneata* (Eoc.-Ala.), 2(2): 35; *ovata* (Mioc.-Calif.), 2(3): 55, 7, 1; *ovula* (Mioc.-Calif.), 2(3): 55, 7, 2; *pseudotorta* (Mioc.-Calif.), 2(3): 55, 7, 3; *rostrata* (Tert.-Venezuela), 5(4): 94, 13, 32; *trinitatensis* (Cret.-Trinidad), 4(4): 102, 14, 12.

Buliminella alabamensis (Eoc.-Ala.), 2(2): 32, 4, 8; *apiculata* (Olig.-Miss.), 5(2): 44, 7, 6, 7; *basistriata* (Eoc.-Trinidad), 5(1): 11, 2, 17-19; *brevior* (Mioc.-Calif.), 1(2): 33, 5, 14; 2(3): 55; *californica* (Mioc.-Calif.), 2(3): 55; *curta* (Mioc.-Calif.), 1(2): 33, 5, 13; 2(3): 55; *elegantissima* (Rec.-Pac.), 1(2): 40, 6, 5; *obtusata* (Olig.-Miss.), 5(2): 42, 7, 8; *subfusiformis* (Mioc.-Calif.), 1(2): 33, 5, 12; 2(3): 55; *subteres*, var. *angusta* (Olig.-Miss.), 5(2): 42, 7, 4.

Calcitornella, 4(2): 45; *elongata* (Penn.-Texas), 4(2): 47, 6, 5; *heathi* (Penn.-Texas), 4(2): 48, 6, 8.

Calcivertella, 4(2): 48; *adherens* (Penn.-Texas), 4(2): 48, 6, 7. [California]. A Cretaceous Cyclammina from California, 4(3): 70. A New Plectofrondicularia from the Pliocene of California, 2(2): 39. Apertural Characters in Cristellaria with Description of a New Species, 1(1): 24. Foraminifera of the Typical Monterey of California, 2(3): 53. Miocene Species of Nonionina

from California, 1(4) : 89. Pliocene Lagenas from California, 5(3) : 67. Siphogenerina hughesi, a New Species from California, 1(2) : 36. Some Later Tertiary Cassidulinas from California, 1(1) : 11. Some Pliocene Bolivinas from California, 2(2) : 40. Some Textulariidae from the Miocene of California, 1(2) : 29. Three New Species of Siphogenerina from the Miocene of California, 1(1) : 2.

Cassidulina, Notes on the Genus, 1(3) : 51.

Cassidulina, 1(3) : 51; *alata* (Mioc.-Italy), 1(3) : 55, 9, 17, 18; *bradyi* (Rec.-Atl.), 1(3) : 52, 8, 3-5; var. *attenuata* (Rec.-Pac.), 1(3) : 53, 8, 16; var. *elongata* (Rec.-Medit.), 1(3) : 53, 8, 14, 15; var. *stenostegica* (Rec.-Atl.), 1(3) : 52, 8, 6; *braziliensis* (Rec.-Atl.), 1(3) : 52, 8, 9, 10; *calabra* (Mioc.-Italy), 1(3) : 55, 9, 14-16; *californica* (Plioc.-Calif.), 1(1) : 12, 2, 1; 1(2) : 40; 1(3) : 54, 9, 5, 6; *charlottensis* (Rec.-Pac.), 1(2) : 41, 6, 6, 7; 1(3) : 53, 8, 17, 18; *clavata* (Rec.-Pac.), 1(3) : 53, 8, 19; *corbyi* (Plioc.-Calif.), 1(1) : 14, 2, 3; 1(3) : 55, 9, 9, 10; *crassa* (Rec.), 1(3) : 54, 8, 37-39; 2(3) : 56, 7, 4; (Tert.-Venezuela), 5(4) : 100, 14, 10; *decorata* (Rec.-Pac.), 1(3) : 53, 8, 20-22; *delicatula* (Tert.-Venezuela), 5(4) : 101, 14, 12; *elegans* (Rec.-Pac.), 1(3) : 53, 8, 23-25; *elegantissima* (Rec.-Pac.), 1(2) : 37, 7, 5; 1(3) : 53, 8, 26, 27; *favus* (Rec.-Pac.), 2(3) : 70; *globosa* (Eoc.-Hungary), 1(3) : 56, 9, 25, 26; *globulosa* (Mioc.-Bavaria), 1(3) : 55; *inexculpta* (Eoc.-Hungary), 1(3) : 56, 9, 27, 28; *laevigata*, 1(2) : 52, 8, 1, 2; var. *carinata* (Rec.-Atl.), 1(3) : 52, 8, 11, 12; var. *tumida* (Rec.-Antarctic), 1(3) : 54, 8, 40-42; *limbata* (Tert.-Calif.), 1(1) : 12, 2, 2; 1(3) : 55, 9, 7, 8; 2(3) : 56; *margareta* (Olig.-Austria), 1(3) : 56, 9, 29, 30; *mexicana* (Rec.-Atl.), 1(3) : 52, 8, 13; *murrhyna* (Plioc.-Pac.), 1(3) : 55, 9, 11; *oblonga* (Mioc.-Austria), 1(3) : 55, 9, 19-22; *obtus* (Rec.-Atl.), 1(3) : 55, 9, 19-22; *orientale* (Rec.-Pac.), 1(2) : 37, 7, 6; 1(3) : 53, 8, 28, 29; *pacifica* (Rec.-Pac.), 1(3) : 53, 9, 14-16; *parkeriana* (Rec.-Pac.), 1(2) : 38, 7, 7; 1(3) : 53, 8, 30, 33; *pulchella* (Plioc.-Calif.), 1(1) : 13, 2, 6; (Mioc.-Calif.), 1(2) : 34, 5, 16; 2(3) : 56; (Rec.-Pac.), 1(2) : 40; 1(3) : 54, 8, 34-36; *punctata* (Mioc.-Austria), 1(3) : 56, 9, 23, 24; *pupa* (Rec.-Antarctic), 1(3) : 54; *sicula* (Pleistoc.-Italy), 1(3) : 54, 8, 46, 47; *spinifera* (Tert.-Trinidad), 5(1) : 17, 3, 1; *subglobosa* (Rec.), 1(3) : 54, 8, 48-50; (Tert.-Venezuela), 5(4) : 100, 14, 11; var. *quadrata* (Pleistoc.-Calif.), 1(1) : 15, 2, 7; 1(3) : 54, 9, 1, 2; var. *tuberculata* (Rec.-Antarctic), 1(3) : 54, 8, 43-45; *tortuosa* (Tert.-Calif.), 1(1) : 14, 2, 4; 1(3) : 55, 9, 12, 13.

Rec.-Pac.), 1(2) : 40; *translucens* (Tert.-Calif.), 1(1) : 15, 2, 3; 1(3) : 54, 9, 3, 4.

sidulinoides, 3(1) : 84.

atobulimina, Some Notes on the Genus, 3(4) : 171.

atobulimina, 3(4) : 171; *alazanensis* (Alazan-Mexico), 3(4) : 74, 29, 5; 30, 3-5; *contraria* (Olig.-Germany), 3(4) : 175, 29, 3; *cretacea* (Cret.-Texas), 3(4) : 173, 29, 1; 30, 11; *dehiscens* (Mioc.-Australia), 3(4) : 176, 29, 7; *evoluta* (Eoc.-Trinidad), 3(1) : 16, 3, 17; *eximia* (Eoc.), 3(4) : 174, 29, 3, 4; 30, 12-16; *guerii* (Mioc.-Austria), 3(4) : 175, 29, 8; 30, 1, 2; var. *australis* (Mioc.-Australia), 3(4) : 176, 29, 10; 30, 6-10; *pacifica* (Rec.-Pac.), 3(4) : 176, 29, 9; *perplexa* (Eoc.-Texas), 3(4) : 173, 29, 2.

lostomella and Related Genera, The Genus, 1(4) : 73.

lostomella, 1(4) : 73; *cylindroides* (Olig.-Germany), 1(4) : 76, 1, 14, 15; *czizeki* (Mioc.-Austria), 1(4) : 74, 11, 2; *grandis* (Rec.-Pac.), 1(4) : 75, 11, 12; *millettii* (Rec.-Pac.), 1(4) : 76, 1, 13; *oolina* (Plioc.-Sicily), 1(4) : 74, 11, 3-10; *ovoidea* (Mioc.-Austria), 1(4) : 74, 11, 1; *tenuis* (Olig.-Germany), 1(4) : 76, 1, 16.

lostomellina, 1(4) : 78; *fimbriata* (Rec.-Pac.), 1(4) : 78, 11, 22.

lostomelloides, 1(4) : 77; *cyclostomata* (Cret.-Hungary), 1(4) : 78, 11, 19; *eocenica* (Eoc.-Texas), 1(4) : 78, 11, 20; *eximia* (Eoc.-Europe), 1(4) : 77, 11, 18; *oviformis* (Eoc.-England), 1(4) : 77, 11, 17, 21.

micidella, 3(1) : 93.

Classification of the Foraminifera, An Outline of a, 3(1) : 1.

bulina amorpha (Cret.-Trinidad), 4(4) : 93, 13, 9; *bramlettei* (Tert.-Ecuador), 5(4) : 81, 12, 11; *curta* (Tert.-Venezuela), 5(4) : 81, 12, 10; *compressa* (Cret.-Texas), 4(3) : 61, 8, 1, 2; *cyclostomata* (Tert.-Ecuador and Venezuela), 5(4) : 80, 12, 8, 9; *trilatera* (Cret.-Trinidad), 4(4) : 93, 13, 8; var. *aspera* (Cret.-Trinidad), 4(4) : 93, 13, 5.

nicospirillina, 3(1) : 73.

rnuspira, 3(4) : 188; cf. *carinata* (Tert.-Venezuela), 5(4) : 82; *cretacea* (Cret.-Mexico), 2(1) : 24, 3, 3; *involvens* (Rec.-Pac.), 1(2) : 44; *thompsoni* (Penn.-Texas), 4(2) : 44, 5, 10.

rnuspiramia, 4(1) : 4; *antillarum* (Rec.-Atl.), 4(1) : 4, 3, 1.

rnuspirella, 4(1) : 4; *diffusa* (Rec.-Atl.), 4(1) : 4, 1, 14.

rnuspiroides, 4(1) : 3; *striolata* (Rec.-Atl.), 4(1) : 3, 1, 13.

[retaceous]. A Cretaceous Cyclammina from California, 4(3) : 70. American Upper Cretaceous Species of *Bolivina* and Related Species, 2(4) : 85. A New Cretaceous *Uvigerina* from

- Louisiana, 1(1): 1. A New Foraminiferal Genus from Upper Cretaceous, 4(1): 12. A Peculiar Clavulina from Upper Cretaceous of Texas, 4(3): 61. Cretaceous Foraminifera from Trinidad, 4(4): 85. Eouvigerina, A New Genus from the Upper Cretaceous, 2(1): 3. Frankeina, A New Genus Arenaceous Foraminifera, 5(3): 61. Kyphopyxa, A New Genus from the Cretaceous of Texas, 5(1): 1. Siphonogenera plummeri, A Species from the Upper Cretaceous of Texas, 2(1): 15. Some Arenaceous Foraminifera from the Upper Cretaceous of Texas, 2(4): 81. Some Foraminifera from Mendez Shale of Eastern Mexico, 2(1): 16. Some New Foraminifera from the Velasco Shale of Mexico, 1(1): 18.
- Cribrobulimina**, 2(4): 80; *mixta* (Rec.-Pac.), 2(4): 80, 11, 1-2.
- Cribrostomoides trinitatensis** (Cret.-Trinidad), 4(4): 91, 12.
- Cristellaria beali** (Mioc.-Calif.), 1(1): 25, 4, 6-13; 2(3): *gutticostata*, var. *cocoaensis* (Eoc.-Ala.), 1(3): 67, 10, *miocenica* (Mioc.-Calif.), 2(3): 57, 7, 8.
- Cyclammina schencki** (Cret.-Calif.), 4(3): 70, 9, 5.
- Cyclocibicides**, 3(1): 93.
- Cycloloculina in the Western Hemisphere, 5(1): 4.
- Cycloloculina**, 3(4): 190; *jarvisi* (Eoc.-Trinidad), 5(1): 5, 1, 8.
- Cymbaloporella**, 3(1): 81.
- Cymbaloporetta**, 4(1): 7.
- Dendritina**, 3(4): 189.
- Dentalina acuta** (Tert.-Venezuela), 5(4): 85, 12, 22; cf. *adolphi* (Cret.-Trinidad), 4(4): 97, 14, 6; (Tert.-Venezuela), 5(4): 13, 3, 4; *consobrina* (Tert.-Venezuela), 5(4): 86, 12, 27; *inornata* (Tert.-Venezuela), 5(4): 85, 12, 21; *pauperata* (Tert.-Venezuela), 5(4): 85, 12, 23, 24; *spinosa* (Tert.-Venezuela), 5(4): 86, 13, 7, 8.
- Depratella, and Its Relation to Endothyra, A New Genus, 4(1): 1.
- Depratella**, 4(1): 9.
- Dimorphina**, 4(1): 16.
- Discorbis charlottensis** (Rec.-Pac.), 1(2): 42, 7, 2; **columbica** (Rec.-Pac.), 1(2): 43, 6, 13; **ornatissima** (Rec.-Pac.), 1(2): 6, 11, 12; **vesicularis** (Eoc.-France), 3(3): 141, 24, 1.
- [Economic]. Foraminifera as an Original Source of Petroleum, 1(2): 48. Some Phases of Correlation by Means of the Foraminifera, 2(3): 71. The Significance of Relative Measurements in the Study of Foraminifera, 2(4): 92.
- Ellipsoidina**, 3(4): 190.
- Ellipsoglandulina exponens** (Cret.-Trinidad), 4(4): 103, 14,

lipsolagena sp. (Plioc.-Calif.), 5(3) : 72, 11, 20.
lipsonodosaria subnodosa (Cret.-Trinidad), 4(4) : 102, 14, 15, 16; *verneuili* (Tert.-Venezuela), 5(4) : 96, 14, 1-3; var. *paucistriata* (Tert.-Ecuador), 5(4) : 97, 14, 4, 5.
lipsopleurostomella curta (Cret.-Trinidad), 4(4) : 102, 14, 18, 19; *pleurostomelloides* (Cret.-Mexico), 2(1) : 18, 2, 6.
phidium crispum (Rec.-Medit.), 5(1) : 20, 4, 3, 4; *macellum* (Rec.-Medit.), 5(1) : 18, 4, 1, 2; *striato-punctatum* (Rec.-Red Sea), 5(1) : 19, 4, 5, 6.
pyrodothyra bowmanni (Penn.-Michigan), 3(2) : 110, 22, 8.
pyrosolenia, 3(4) : 190; *staphyllearia* (Tert.-Venezuela), 5(4) : 96, 13, 40.
 Eocene]. A Peculiar Frondicularia from Mexico and Trinidad, 1(4) : 88. A Species of Siphoninella in the Tertiary of America, 3(2) : 120. Additional Foraminifera from the Eocene of Alabama, 4(3) : 73. Cycloloculina in the Western Hemisphere, 5(1) : 4. Eocene Foraminifera from the Cocoa Sand of Alabama, 1(3) : 65. New Foraminifera from the Upper Eocene of Mexico, 1(1) : 4. Some New Foraminifera from the Upper Eocene of the Southeastern Coastal Plain of the United States, 2(2) : 29.
pyrogerina, 2(1) : 4; *americana* (Cret.-Texas), 2(1) : 4, 1, 1; *gracilis* (Cret.-Texas), 2(1) : 5, 1, 2.
pyropistomella, 4(1) : 6; *rimosa*, 4(1) : 6, 1, 9.
pyropistomina, 3(4) : 181; *elegans*, 3(4) : 182, 31, 32.
pyropistomina elegans (d'Orbigny) and *E. partschiana* (d'Orbigny), 3(4) : 180.
pyropistomides cocoaensis (Eoc.-Ala.), 4(3) : 73, 10, 2; *umbonata* (Tert.-Venezuela), 5(4) : 98, 14, 8.
 Europe]. A New Uvigerina from the Vienna Basin, 1(1) : 10.
pyropistomina d'Orbigny, On, 4(3) : 80.
pyropistomina, 4(3) : 80.
pyropistomina, 4(1) : 1; *alexanderi* (L. Cret.-Texas), 4(1) : 1, 1, 3, 4.
pyropistomina, 3(4) : 189; *semireticulata* (Cret.-Trinidad), 4(4) : 98, 13, 14.
 Florida]. An American *Virgulina* Related to *V. pertusa* Reuss, 5(3) : 53.
pyropistomina, 5(3) : 61; *goodlandensis* (Cret.-Texas), 5(3) : 62, 10, 1, 2; *taylorensis* (Cret.-Texas), 5(3) : 63, 10, 3.
pyropistomina archiaciana (Cret.-Mexico), 2(1) : 21, 3, 4; *baudouiniana* (Cret.-Mexico), 2(1) : 21, 3, 5; *elongata* (Cret.-Trinidad), 4(4) : 98, 14, 1; *foliacea* (Mioc.-Calif.), 2(3) : 57, 7, 9;

- mexicana* (Alazan-Mexico), 1(4) : 88, 13, 5; *tenuissima* (Alazan-Mexico), 3(2) : 111, 22, 11.
- Gaudryina and Heterostomella, Fistulose Species of, 4(4) : 10.
- Gaudryina disjuncta* (Eoc.-Trinidad), 5(1) : 6, 2, 7; *filiformis* (Cret.-Trinidad), 4(4) : 92, 13, 2; *gardnerae* (Eoc.-Ala.), 2(2) : 33, 5, 2; *indentata* (Cret.-Trinidad), 4(4) : 92, 13, 7; *jacksonensis* (Eoc.-So. Car.), 2(2) : 33, 5, 1; *laevigata*, var. *pyramidalis* (Cret.-Trinidad), 4(4) : 92, 13, 6; *retusa* (Cret.-Trinidad), 4(4) : 92, 13, 3, 4; *siphonifera* (Rec.-Pac.), 4(4) : 109, 16, 1; *stephensoni* (Cret.-Texas), 4(4) : 108, 16, 6-8; *trinitatis* (Tert.-Ecuador), 5(4) : 80, 12, 7; *velascoensis* (Cret.-Mexico), 1(1) : 20, 3, 7.
- Geinitzina ciscoensis* (Penn.-Texas), 4(3) : 63, 8, 6.
- Glandulina*, 3(4) : 189; *laevigata* (Tert.-Venezuela), 5(4) : 13, 17.
- Globigerina mexicana* (Eoc.-Mexico), 1(1) : 6, 1, 8; *topilensis* (Eoc.-Mexico), 1(1) : 7, 1, 9; *velascoensis* (Cret.-Mexico), 1(1) : 19, 3, 6.
- Globigerinella*, 3(1) : 87.
- Globigerinoides*, 3(1) : 87.
- Globivalvulina biserialis* (Penn.-Texas), 4(3) : 64, 8, 7; *trilineata* (Penn.-Texas), 4(3) : 65, 8, 8.
- Globobulimina*, 3(1) : 67; *pacifica* (Rec.-Pac.), 3(1) : 67, 14, 1.
- Globorotalia*, 3(1) : 91; *canariensis* (Tert.-Venezuela), 5(4) : 14, 14; *cocoaensis* (Eoc.-Ala.), 4(3) : 75, 10, 3; *lehneri* (Cret.-Trinidad), 5(1) : 17, 3, 16; *spinulosa* (Alazan-Mexico), 3(2) : 114, 23, 4.
- Globotruncana*, 3(1) : 91; *calcarata* (Cret.-Texas), 3(2) : 110, 10; *canaliculata* (Cret.-Texas), 3(2) : 116, 23, 11.
- Globulina*, 3(4) : 189.
- Glomospira charoides*, var. *corona* (Cret.-Trinidad), 4(4) : 88, 9-11; *duplex* (Penn.-Texas), 3(3) : 147, 26, 6; *gordialis* (Cret.-Trinidad), 4(4) : 87, 12, 7, 8; *pusilla* (Penn.-Michigan), 3(3) : 108, 22, 1; *spinosa* (Penn.-Texas), 4(2) : 39, 4, 14; *umbilicatus* (Penn.-Texas), 3(3) : 148, 26, 7, 8.
- Glyphostomella*, 4(2) : 53; *triloculina* (Penn.-Texas), 4(2) : 111-13; 7, 1.
- Gonatosphaera prolata* (Tert.-Venezuela), 5(4) : 97, 13, 41.
- Guembelina*, 3(4) : 190; *excolata* (Cret.-Mexico), 2(1) : 20, 2.
- Guttulina*, 4(1) : 14; *irregularis* (Tert.-Venezuela), 5(4) : 88, 15, 16; *lactea* (Rec.-Atl.), 5(2) : 36, 6, 6-10; *orientalis* (Eoc.-Japan), 4(1) : 15, 2, 1.

Gyroidina, 3(4) : 190; *soldanii* (Tert.-Venezuela), 5(4) : 98, 14, 6, 7.

Hantkenina alabamensis (Alazan-Mexico), 1(1) : 7, 1, 11; (Eoc.-Ala.), 1(3) : 68; var. **primitiva** (Eoc.-Trinidad), 5(1) : 16, 3, 2, 3; *brevispina* (Eoc.-Mexico), 1(1) : 7, 1, 10; **lehneri** (Eoc.-Trinidad), 5(1) : 16, 3, 8.

Haplophragmium taylorensis (Cret.-Texas), 5(3) : 64, 10, 4, 5.

Haplophragmoides advena (Rec.-Pac.), 1(2) : 38, 6, 1; **calcula** (Cret.-Texas), 2(4) : 83, 10, 5; **columbiensis** (Rec.-Pac.), 1(2) : 39, 6, 2; *coronata* (Cret.-Trinidad), 4(4) : 90, 12, 17; *emaciatum* (Tert.-Venezuela), 5(4) : 78, 12, 3; *excavata* (Cret.-Texas), 2(4) : 82, 10, 3; **glabra** (Cret.-Texas), 2(4) : 83, 10, 6; **rugosa** (Cret.-Texas), 2(4) : 83, 10, 4; cf. *subglobosum* (Cret.-Trinidad), 4(4) : 91, 12, 13; sp(?) (Cret.-Trinidad), 4(4) : 91, 12, 16.

Hastigerinella, 3(1) : 87.

Hemigordius calcarea (Permian-Texas), 4(2) : 44, 6, 1, 2; **harltoni** (Penn.-Texas), 4(2) : 43, 5, 8, 9; **liratus** (Penn.-Texas), 4(2) : 44, 5, 11, 12.

Heterohelix, 3(4) : 190.

Heterostomella, *Fistulose Species of Gaudryina* and, 4(4) : 107.

Heterostomella foveolata (Cret.), 4(4) : 111, 16, 9-12.

Hormosina globulifera (Cret.-Trinidad), 4(4) : 86, 12, 3.

Howchinia, 3(1) : 42.

Hyperammia bulbosa (Penn.-Michigan), 3(2) : 109, 22, 7; *elongata* (Cret.-Trinidad), 4(4) : 86, 12, 1; **glabra** (Penn.-Texas), 3(3) : 146, 26, 1; *spinescens* (Penn.-Texas), 4(2) : 35, 4, 1.

Hyperamminella, 4(2) : 36; **elegans** (Penn.-Texas), 4(2) : 36, 4, 3, 4; **minuta** (Penn.-Texas), 4(2) : 37, 4, 8, 9; **protea** (Penn.-Texas), 4(2) : 36, 4, 5, 6.

Hyperamminoides, A New Name for *Hyperamminella*, 4(4) : 112.

Hyperamminoides, 4(4) : 112.

Kyphopyxa, A New Genus from the Cretaceous of Texas, 5(1) : 1.

Kyphopyxa, 5(1) : 1; *christneri* (Cret.-Texas), 5(1) : 1, 1, 1-7.

Lagena angelina (Plioc.-Calif.), 5(3) : 71, 11, 10; **byramensis** (Olig.-Miss.), 5(2) : 41, 7, 3; *clavata* (Plioc.-Calif.), 5(3) : 68, 11, 3; *costata* (Plioc.-Calif.), 5(3) : 70, 11, 9; var. *amphora* (Plioc.-Calif.), 5(3) : 70, 11, 11, 12; *clypeata* (Plioc.-Calif.), 5(3) : 72, 11, 19; *elongata* (Plioc.-Calif.), 5(3) : 67, 11, 1; *gracilis* (Plioc.-Calif.), 5(3) : 67, 11, 2; (Tert.-Venezuela), 5(4) : 88, 13, 11; *hexagona* (Plioc.-Calif.), 5(3) : 72, 11, 18; var. *scalariformis* (Plioc.-Calif.), 5(3) : 72, 11, 17; *hispida*

- (Plioc.-Calif.), 5(3): 71, 11, 13; *lacuno-costata* (Eoc.-Trinidad), 5(1): 8, 2, 8, 9; *lamellata* (Plioc.-Calif.), 5(3): 71, 11, 14; *marginata* (Plioc.-Calif.), 5(3): 71, 11, 15; *orbignyana* (Rec.-Pac.), 1(2): 41, 6, 8; var. *elliptica* (Plioc.-Calif.), 5(3): 71, 11, 16; *pulcherrima* (Tert.-Trinidad), 5(1): 8, 2, 10; *sacculus* (Plioc.-Calif.), 5(3): 72, 11, 21; *striata* var. *strumosa* (Plioc.-Calif.), 5(3): 70, 11, 6; *substriata* (Plioc.-Calif.), 5(3): 68, 11, 4; *sulcata* (Plioc.-Calif.), 5(3): 70, 11, 5; *williamsoni* (Plioc.-Calif.), 5(3): 70, 11, 7, 8; sp(?) (Tert.-Venezuela), 5(4): 88, 12, 30.
- Lamarckina* and Its American Species, The Genus, 2(1): 7.
- Lamarckina*, 2(1): 7; *cristellaroides* (Eoc.-France), 2(1): 10; *erinacea* (Mioc.-Hungary), 2(1): 12; *glabrata* (Olig.-Miss.), 2(1): 11, 1, 6; *halkyardi* (Eoc.-France), 2(1): 11; *marylandica* (Eoc.-Md.), 2(1): 9, 1, 5; var. *claibornensis* (Eoc.-Ala.), 2(1): 10; var. *yeguaensis* (Eoc.-Texas), 2(1): 10; *ocalana* (Eoc.-Fla.), 2(1): 11, 1, 4; *ripleyensis* (Cret.-Miss.), 2(1): 8; *rugulosa* (Eoc.-Texas), 2(1): 8, 3, 6; *scabra* (Rec.), 2(1): 13; *umbilicata* (Mioc.-Australia), 2(1): 12; *ventricosa* (Rec.-Atl.), 2(1): 12; *wilcoxensis* (Eoc.-Ala.), 2(1): 9, 1, 3.
- Lenticulina grata* (Cret.-Trinidad), 4(4): 96, 14, 3; *nuda* (Cret.-Trinidad), 4(4): 96, 14, 2; *rotulata* (Cret.-France), 3(3): 142, 28, 7.
- Lituola nautiloidea* (Eoc.-France), 3(3): 142, 28, 8, 9; *taylorensis* (Cret.-Texas), 5(3): 66, 10, 7.
- Lituotuba lituiformis* (Cret.-Trinidad), 4(4): 90, 12, 15.
- Louisiana, A New Cretaceous Uvigerina from, 1(1): 1.
- Loxostomum amygdalaeformis*, var. *delicata* (Olig.-Miss.), 5(2): 44, 8, 2.
- Marginulina, Mexican Species of, 1(3): 61.
- Marginulina bullata* (Cret.-Trinidad), 4(4): 96, 14, 7, 8; *cocoaensis* (Eoc.-Ala.), 1(3): 67, 10, 9, 10; *mexicana* (Olig.-Mexico), 1(3): 61, 10, 1; *pulchra* (Olig.-Mexico), 1(3): 61, 10, 2; *subbullata* (Olig.-Mexico), 1(3): 62, 10, 3; (Tert.-Venezuela), 5(4): 85, 12, 20.
- Massilina decorata* (Olig.-Miss.), 5(2): 40, 7, 1.
- [Mexico]. A Peculiar Frondicularia from Mexico and Trinidad, 1(4): 88. Mexican Species of Marginulina, 1(3): 61. New and Interesting Foraminifera from Mexico and Texas, 3(2): 111. New Foraminifera from the Upper Eocene of Mexico, 1(1): 4. Some Foraminifera from the Mendez Shale of Eastern Mexico, 2(1): 16. Some Fossil Bolivinas from Mexico, 1(4): 81. Some New Foraminifera from the Velasco Shale of Mexico, 1(1): 18.

Michigan, Pennsylvanian Foraminifera from, 3(2) : 107.

Mimosina, 3(4) : 190.

[Miocene]. An American *Virgulina* Related to *V. pertusa* Reuss, 5(3) : 53. A New *Uvigerina* from the Vienna Basin, 1(1) : 10. Apertural Characters in *Cristellaria* with Descriptions of a New Species, 1(1) : 24. Foraminifera of the Typical Monterey of California, 2(3) : 53. Miocene Species of *Nonionina* from California, 1(4) : 89. *Siphogenerina hughesi*, a New Species from California, 1(2) : 36. Some Textulariidae from the Miocene of California, 1(2) : 29. Three New Species of *Siphogenerina* from the Miocene of California, 1(1) : 2.

[Mississippi]. Notes on the Foraminiferal Fauna of the Byram Marl, 5(2) : 40.

Masonella, 3(4) : 188.

Monalysidium, 3(4) : 189.

Mooreinella, 4(2) : 50; *biserialis* (Penn.-Texas), 4(2) : 50, 6, 9, 10.

Neobulimina, 4(1) : 12; *canadensis* (Cret.-Canada), 4(1) : 13, 1, 1, 2.

Neocribrella, 4(1) : 6; *globigerinoides* (Eoc.-France), 4(1) : 6, 1, 6, 7.

Nodogenerina, 2(4) : 79; *bradyi* (Rec.-Pac.), 2(4) : 79.

Nodomorphina, 2(4) : 80.

Nodosaria acuminata (Tert.-Venezuela), 5(4) : 86, 13, 5, 6; *arundinea* (Tert.-Ecuador), 5(4) : 86, 13, 1, 2; *brevitesta* (Cret.-Trinidad), 4(4) : 97, 13, 10; *cocoaensis* (Eoc.-Ala.), 1(3) : 66, 10, 5, 6; *concinna* (Cret.-Trinidad), 4(4) : 97, 14, 5, 11; *fissicostata* (Eoc.-Ala.), 1(3) : 66, 10, 8; *koina* (Mioc.-Calif.), 2(3) : 56, 7, 6; var. *hughesi* (Mioc.-Calif.), 2(3) : 57, 7, 7; *latejugata* (Eoc.-Ala.), 1(3) : 66, 10, 7; *limonensis* (Cret.-Mexico), 1(1) : 21, 3, 4; *longiscata* (Tert.-Venezuela), 5(4) : 86, 12, 25, 26; cf. *mareki* (Cret.-Trinidad), 4(4) : 97, 14, 4; *mexicana* (Eoc.-Mexico), 1(1) : 5, 1, 3, 4; *obliqua* (Mioc.-Calif.), 2(3) : 56, 7, 5; *obscura* (Cret.-Mexico), 2(1) : 20, 2, 8; *schlichti* (Tert.-Venezuela), 5(4) : 87, 13, 9, 10; *velascoensis* (Cret.-Trinidad), 4(4) : 97, 13, 15, 16; *vertebralis* (Tert.-Venezuela), 5(4) : 86.

Nodosinella, 3(4) : 188; *arenata* (Penn.-Texas), 3(3) : 147, 26, 2, 3; *glabra* (Penn.-Texas), 3(3) : 147, 26, 4, 5.

Nonion affinis (Tert.-Venezuela), 5(4) : 89, 13, 24; *pompilioides* (Tert.-Venezuela), 5(4) : 89, 13, 25.

Nonionina auris (Rec.-Pac.), 1(2) : 44, 7, 3; (Mioc.-Calif.), 1(4) : 91, 13, 14; *costifera* (Mioc.-Calif.), 1(4) : 90, 13, 2; 2(3) : 65;

- incisa* (Mioc.-Calif.), 1(4) : 90, 13, 3; 2(3) : 65; *medio-costata* (Mioc.-Calif.), 1(4) : 89, 13, 1; 2(3) : 65.
- Nonionella*, 2(3) : 64; *miocenica* (Mioc.-Calif.), 2(3) : 64.
- Normanina*, 4(1) : 7.
- Nouria*, 3(4) : 189.
- [Oligocene]. Notes on the Foraminiferal Fauna of the Byram Marl, 5(2) : 40.
- Orthovertella*, 4(2) : 45; *protea* (Penn.-Texas), 4(2) : 45, 6, 3,
- [Palaeozoic]. A New Genus, *Depratella*, and its Relation to *Endothyra*, 4(1) : 9. Arenaceous Palaeozoic Foraminifera from Texas, 3(3) : 146.
- Patellina protea* (Permian-Texas), 4(2) : 54, 7, 8-10.
- Patellinella*, 4(1) : 5; *inconspicua*, 4(1) : 5, 1, 8.
- [Pennsylvanian]. Additional Cisco Foraminifera from Texas, 4(3) : 62. Pennsylvanian Foraminifera from Michigan, 3(2) : 107. Some Foraminifera from the Pennsylvanian and Permian of Texas, 4(2) : 31. The Microspheric and Megalospheric Forms of *Apterrinella grahamensis*, 4(3) : 68.
- [Permian]. Some Foraminifera from the Pennsylvanian and Permian of Texas, 4(2) : 31.
- Planoglobulina*, 2(4) : 77.
- Planorbulinella*, 3(1) : 96.
- Planorbulinoides*, 4(1) : 6; *retinaculata* (Rec.-Pac.), 4(1) : 6, 1.
- Planularia clara* (Tert.-Trinidad), 5(1) : 7, 2, 14, 15.
- Planulina ariminensis* d'Orbigny and *P. wuellerstorfi* (Schwager), 5(4) : 102.
- Planulina ariminensis* (Rec.-Medit.), 5(4) : 102, 15, 3, 4; *coarctata* (Eoc.-Ala.), 4(3) : 76, 10, 1; *mexicana* (Alazan-Mexico), 3(2) : 113, 23, 5; *wuellerstorfi* (Plioc.-Pac.), 5(4) : 104, 15, 17.
- Plectofrondicularia alazanensis* (Alazan-Mexico), 3(2) : 113, 12; *californica* (Mioc.-Calif.), 2(2) : 39, 6, 9-11; (Tert.-Venezuela), 5(4) : 90, 13, 18, 19; *miocenica* (Mioc.-Calif.), 2(3) : 7, 10, 11; 8, 11, 12; (Tert.-Venezuela), 5(4) : 92, 13, 20; *mormonensis* (Tert.-Ecuador), 5(4) : 92, 13, 23; *paucicostata* (Eoc.-Trinidad), 5(1) : 10, 2, 11-13; *spinifera* (Eoc.-Trinidad), 5(1) : 2, 1-3; *trilineata* (Alazan-Mexico), 3(2) : 112, 23, 2; *trinidadensis* (Eoc.-Trinidad), 5(1) : 11, 2, 16; *vaughani* (Alazan-Mexico), 3(2) : 112, 23, 3; (Tert.-Venezuela), 5(4) : 92, 21, 22.
- Pleurostomella*, Notes on the Genus, 3(2) : 128. Additional Notes on the Genus *Pleurostomella*, 3(3) : 156.
- Pleurostomella*, 3(2) : 128; *acuminata* (Rec.-Atl.), 3(2) : 129,

1; *acuta* (Eoc.-Hungary), 3(2): 130, 25, 14; var. *buliminiformis* (Plioc.-Italy), 3(3): 156, 28, 5; *alazanensis* (Alazan-Mexico), 1(1): 5, 1, 2; 3(2): 130, 25, 21; *alternans* (Plioc.-Ind. Pac.), 3(2): 129, 25, 7, 8, 28; (Tert.-Venezuela), 5(4): 96, 13, 42, 43; var. *hians* (Mioc.-Austria), 3(2): 130, 25, 12; var. *moravica* (Mioc.-Austria), 3(2): 130, 25, 11; var. *parvifnita* (Mioc.-Austria), 3(2): 130, 25, 10; var. *telostoma* (Mioc.-Austria), 3(2): 130, 25, 9; *antiqua* (Permo-Carb.-Australia), 3(2): 132, 25, 27; *barroisi* (Cret.-France), 3(2): 131, 25, 24; *bellardi* (Eoc.-Hungary), 3(2): 131; 3(3): 156, 28, 1; *brevis* (Plioc.-Ind. Pac.), 3(2): 129, 25, 6; *clavata* (Cret.-Mexico), 3(2): 132, 25, 19; *contorta* (Rec.-Pac.), 3(2): 129, 25, 3; *eocaena* (Eoc.-Bavaria), 3(2): 130, 25, 15; *fusiformis* (Cret.-Germany), 3(2): 132, 25, 21; *globulifera* (Cret.-Germany), 3(2): 132; *incrassata* (Eoc.-Hungary), 3(2): 131; 3(3): 156, 28, 2, 3; *jacksonensis* (Eoc.-Texas), 3(2): 131, 25, 17; *jurassica* (Jurassic-Switz.), 3(2): 132, 25, 30-32; *obtusa* (Cret.-France), 3(2): 131, 25, 22; *pleurostomella* (Mioc.-Italy), 3(2): 130, 25, 13; *rapa* (Eoc.-Bavaria), 3(2): 131, 25, 18; var. *recens* (Rec.-Ind. Pac.), 3(3): 156, 28, 6; *reussi* (Cret.-France), 3(2): 132, 25, 25; *sapperi* (Pleist.-Pac.), 3(2): 129, 25, 4; *schuberti* (Rec.-Pac.), 3(2): 133, 25, 5, 29; *spinosa* (Rec.-Pac.), 3(2): 129, 25, 2; *subnodosa* (Cret.-Germany), 3(2): 131, 25, 23; *tenuis* (Eoc.-Hungary), 3(2): 131; 3(3): 156, 28, 4; *torta* (Cret.-Mexico), 2(1): 18, 2, 7; 3(2): 132, 25, 16; *velascoensis* (Cret.-Mexico), 3(2): 132, 25, 20.

[Pliocene]. A New Plectofrondicularia from the Pliocene of California, 2(2): 39. Pliocene Lagenas from California, 5(3): 67. Some Late Tertiary Cassidulinas from California, 1(1): 11. Some Pliocene Bolivinas from California, 2(2): 40.

Plummerinella, 4(2): 49; **complexa** (Penn.-Texas), 4(2): 49, 6, 6. *Polymorphina*, 4(1): 16; *advena* (Olig.-Miss.), 5(2): 41, 7, 5; *burdigalensis* (Mioc.-France), 5(2): 35, 6, 1-3; **charlottensis** (Rec.-Pac.), 1(2): 41, 6, 9; *frondea* (Olig.-Miss.), 5(2): 41; **jacksonensis** (Eoc.-Fla.), 2(2): 35, 5, 5; var. **costifera** (Eoc.-Ala.), 2(2): 35.

Polytaxis, 4(2): 51; **laheei** (Penn.-Texas), 4(2): 51, 7, 7; **multi-loculata** (Penn.-Texas), 4(2): 51.

Problematina, 3(4): 188.

Proroporus, 2(4): 89; *plaita* (Cret.-Texas), 2(4): 89, 12, 7.

Proteonina cervicifera (Penn.-Texas), 4(2): 33, 4, 11; *diffugiiformis* (Cret.-Texas), 2(4): 82, 10, 1; (Tert.-Venezuela), 5(4): 78, 12, 2.

- Psammophis inclusus* (Penn.-Texas), 3(3) : 148, 26, 12; *inversus* (Penn.-Michigan), 3(2) : 109, 22, 4.
- Pseudoglandulina* 5(4) : 87; *gallowayi* (Tert.-Ecuador), 5(4) : 87, 13, 13.
- Pseudopolymorphina*, 4(1) : 15; *hanzawai* (Plioc.-Japan), 4(1) : 16, 1, 15.
- Pseudotextularia acervulinoides* (Cret.-Mexico), 2(1) : 17, 2, 5; *varians* (Cret.-Mexico), 2(1) : 17, 2, 4.
- Pseudouvigerina*, 2(4) : 81; *plummerae* (Cret.-Texas), 3(2) : 115, 23, 8.
- Pulleniatina*, 3(1) : 90.
- Pulvinulina arca* (Cret.-Mexico), 2(1) : 23, 3, 1; var. *contusa* (Cret.-Mexico), 2(1) : 23; *columbiensis* (Rec.-Pac.), 1(2) : 43, 7, 1; *repanda* (Rec.-Pac.), 1(2) : 43; *velascoensis* (Cret.-Mexico), 1(1) : 19, 3, 5.
- Pulvinulinella*, 2(3) : 62; *culter* (Tert.-Venezuela), 5(4) : 100, 14, 13; *exigua* (Tert.-Ecuador), 5(4) : 100; *interrupta* (Alazan-Mexico), 3(2) : 113, 22, 10; *subperuviana* (Mioc.-Calif.), 2(3) : 63, 9, 9.
- Pyrulina gutta* (Plioc.-Italy), 5(2) : 39, 6, 4, 5.
- Pyrulinella*, 4(1) : 16.
- Quinqueloculina seminula* (Rec.-Medit.), 5(3) : 59, 9, 16-18.
- Ramulina*, 4(1) : 19.
- Recent Foraminifera from British Columbia, 1(2) : 38.
- Rectobolivina*, 3(1) : 68.
- Renulina* and *Vertebralina*, The Genera, 3(3) : 158.
- Renulina opercularia* (Eoc.-France), 3(3) : 158, 28, 10.
- Reophax asperus* (Penn.-Texas), 4(2) : 37, 4, 7; *texana* (Cret.-Texas), 2(4) : 82, 10, 2; sp(?) (Cret.-Trinidad), 4(4) : 86, 12, 2.
- Robulus calcar* (Tert.-Venezuela), 5(4) : 84, 12, 18; *clericii* (Tert.-Venezuela), 5(4) : 84, 12, 16, 17; *subaculeata*, var. *glabrata* (Tert.-Venezuela), 5(4) : 85, 12, 19; sp(?) (Tert.-Venezuela), 5(4) : 82, 84.
- Rotalia beccarii* Linné, On, 4(4) : 103.
- Rotalia beccarii* (Rec.-Medit.), 4(4) : 104, 15, 1-7; *trochidiformis* (Eoc.-France), 3(3) : 142, 24, 5-7.
- Rotaliatina*, 1(1) : 4; *mexicana* (Alazan-Mexico), 1(1) : 4, 1, 1.
- Rzehakina*, 3(1) : 31; *epigona*, var. *lata* (Cret.-Trinidad), 4(4) : 93, 13, 11.
- Saracenaria acutaureicularis* (Tert.-Venezuela), 5(4) : 88, 13, 12; *italica* (Tert.-Venezuela), 5(4) : 88, 13, 14.

Sigmoidella, 4(1) : 18; **kagaensis** (Plioc.-Japan), 4(1) : 19, 2, 14;
(Sigmoidina) pacifica (Rec.-Pac.), 4(1) : 19, 2, 13.

Sigmoidina, 4(1) : 18.

Sigmoilina celata (Tert.-Venezuela), 5(4) : 82; **tenuis** (Tert.-Venezuela), 5(4) : 81, 12, 12-14.

Sigmomorpha, 4(1) : 17; **(Sigmomorphina) yokoyamai** (Plioc.-Japan), 4(1) : 18, 1, 16.

Sigmomorphina, 4(1) : 17; **sadoensis** (Plioc.-Japan), 4(1) : 17, 2, 11.

Silicina, 3(4) : 188.

Siphogenerina, 3(4) : 190; **basispinata** (Tert.-Trinidad), 5(1) : 13, 3, 4, 5; **collomi** (Mioc.-Calif.), 1(1) : 2, 4, 3; 2(3) : 59; **hughesi** (Mioc.-Calif.), 1(2) : 36, 7, 4; **kleinpelli** (Mioc.-Calif.), 1(1) : 3, 4, 5; 2(3) : 59; **multicostata** (Tert.-Trinidad), 5(1) : 14, 3, 6; (Tert.-Venezuela), 5(4) : 95, 13, 38; **plummeri** (Cret.-Texas), 2(1) : 15, 1, 7; **raphana**, var. **transversa** (Tert.-Venezuela), 5(4) : 95; **reedi** (Mioc.-Calif.), 1(1) : 3, 4, 4; 2(3) : 59.

Siphogenerinoides, 3(1) : 63; 5(3) : 55; **bramletti** (Cret.-Venezuela), 5(3) : 56, 9, 5, 6; **cretacea** (Cret.-Venezuela), 5(3) : 58, 9, 14, 15; **ewaldi** (Cret.-So. Amer.), 5(3) : 58; **parva** (Cret.-Venezuela), 5(3) : 58, 9, 11-13; **plummeri** (Cret.-Venezuela), 5(3) : 55.

Siphonina tenuicarinata (Tert.-Venezuela), 5(4) : 100, 14, 9.

Siphoninella, 3(1) : 77; **byramensis** (Olig.-Miss.), 5(2) : 46, 8, 5; **claibornensis** (Eoc.-La.), 3(2) : 120, 24, 8-10.

Siphoninoides, 3(1) : 77.

Sorites, 3(4) : 190.

Sphaeroidina variabilis (Tert.-Venezuela), 5(4) : 101, 14, 15.

Sphaeroidinella, 3(1) : 90.

Spirillina spinigera, var. **reducta** (Rec.-Pac.), 1(2) : 42, 6, 10.

Spiroclypeus, 3(4) : 189.

Spiroloculina limbata (Tert.-Venezuela), 5(4) : 82, 12, 15.

Spiroplectammina, 3(1) : 23; **castensis** (Penn.-Texas), 4(3) : 62, 8, 4; **clavata** (Penn.-Texas), 3(3) : 150, 26, 11.

Spiroplectina, 2(4) : 78.

Spiroplectinata, 3(1) : 62.

Spiroplectoides, 2(4) : 77; **clotho** (Cret.-Trinidad), 4(4) : 101, 14, 13, 14; **rosula** (Cret.-Texas), 3(2) : 114, 23, 6, 7.

Spirothamidium, 3(1) : 37.

Sporadogenerina, A Degenerate Foraminiferal Genus, 2(4) : 94.

Sporadogenerina, 2(4) : 95; **flintii** (Rec.-Atl.), 2(4) : 95, 11, 6, 7.

Stacheia, 3(4) : 189.

[Structural]. Apertural Characters in Cristellaria with Descriptions of a New Species, 1(1): 24. Apertural Characters in the Lagenidae, 4(1): 22. The Term "Arenaceous Foraminifera" and Its Meaning, 5(2): 25.

[Taxonomic]. Additional Genera of the Foraminifera, 4(1): 1. An Outline of a Revision of the Polymorphinidae, 4(1): 13. Notes on the Collection of DeFrance, 3(3): 141. On *Elphidium macellum* (Fichtel and Moll), *E. striato-punctatum* (Fichtel and Moll) and *E. crispum* (Linné), 5(1): 18. On *Guttulina lactea* (Walker and Jacob), *Polymorphina burdigalensis* d'Orbigny and *Pyrulina gutta* d'Orbigny, 5(2): 34. On *Quinqueloculina seminula* (Linné), 5(3): 59. Some New Genera of the Foraminifera, 2(4): 77. Some Notes on the Early Foraminiferal Genera Erected before 1808, 3(2): 122. The Designation of Some Genotypes in the Foraminifera, 3(4): 188. The Generic Position of *Pulvinulina favus* H. B. Brady, 2(3): 70. The Work of Fichtel and Moll and of Montfort, 3(4): 168. Trimorphism in the Foraminifera, 3(4): 165.

[Technique]. Photographing Foraminifera, 2(1): 1.

Tetrataxis corona (Penn.-Texas), 4(3): 65, 8, 10; *millsapensis* (Penn.-Texas), 4(2): 50, 7, 5, 6; *multiloculata* (Penn.-Texas), 3(3): 153, 27, 6; *scutella* (Penn.-Texas), 4(3): 65, 8, 9.

[Texas]. A Peculiar Clavulina from the Upper Cretaceous of Texas, 4(3): 61. Additional Cisco Foraminifera from Texas, 4(3): 62. Arenaceous Palaeozoic Foraminifera from Texas, 3(3): 146. *Eouvigerina*, A New Genus from the Upper Cretaceous, 2(1): 3. *Frankeina*, A New Genus of Arenaceous Foraminifera, 5(3): 61. *Kyphopyxa*, A New Genus from the Cretaceous of Texas, 5(1): 1. New and Interesting Foraminifera from Mexico and Texas, 3(2): 111. *Siphogenerina plummeri*, A Species from the Upper Cretaceous of Texas, 2(1): 15. Some Arenaceous Foraminifera from the Taylor Marl of Texas, 5(3): 63. Some Arenaceous Foraminifera from the Pennsylvanian and Permian of Texas, 4(2): 31.

Textularia adalta (Eoc.-Ala.), 2(2): 29, 4, 2; *concinna* (Cret.-Trinidad), 4(4): 91, 13, 1; *exrayensis* (Penn.-Texas), 3(3): 150, 27, 2; *fuscalignensis* (Penn.-Texas), 3(3): 151, 27, 1; *grahamensis* (Penn.-Texas), 3(3): 151, 27, 3; *lirata* (Tert.-Trinidad), 5(1): 6, 2, 4; *mexicana* (Tert.-Venezuela), 5(4): 79, 12, 4; *mississippiensis* (Tert.-Venezuela), 5(4): 79, 12, 5; *ocalana* (Eoc.-Fla.), 2(2): 30, 4, 3; *velascoensis* (Cret.-Mexico), 1(1): 18, 3, 1.

Textulariella, 3(1): 24.

- Thurammina texana* (Penn.-Texas), 4(2) : 33, 4, 2.
- Tolypammina delicatula* (Penn.-Texas), 4(3) : 2, 8, 3.
- Trepeilopsis**, 4(2) : 38; *grandis* (Penn.-Texas), 4(2) : 38, 4, 12, 13.
- Trifarina in the American Eocene and Elsewhere, 1(4) : 86.
- Trifarina*, 1(4) : 86; *bradyi* (Rec.), 1(4) : 86; (Tert.-Venezuela), 5(4) : 96, 13, 39; var. *advena* (Eoc.-Miss.), 1(4) : 87; *reussi* (Rec.), 1(4) : 87.
- Triloculina sculpturata* (Olig.-Miss.), 5(2) : 40, 7, 2.
- Trimosina**, 3(1) : 64.
- [Trinidad]. A Peculiar Frondicularia from Mexico and Trinidad, 1(4) : 88. Cretaceous Foraminifera from Trinidad, 4(4) : 85. *Cycloloculina* in the Western Hemisphere, 5(1) : 4. New Foraminifera from Trinidad, 5(1) : 6.
- Tritaxilina**, Notes on the Genus, 1(3) : 62.
- Tritaxilina*, 1(3) : 63; *atlantica* (Rec.-Atl.), 1(3) : 64; *caperata* (Rec.-Pac.), 1(3) : 63; *mexicana* (Eoc.-Mexico), 1(3) : 64, 10, 4.
- Trochammina arenosa* (Penn.-Texas), 3(3) : 152, 27, 4; **charlottensis** (Rec.-Pac.), 1(2) : 39, 6, 4; *diagonis* (Cret.-Texas), 2(4) : 84, 10, 7; *globigeriniformis* (Cret.-Trinidad), 4(4) : 95, 13, 12; *gyroides* (Cret.-Texas), 2(4) : 84, 10, 8; *pacifica* (Rec.-Pac.), 1(2) : 39, 6, 3; *rudis* (Penn.-Michigan), 3(2) : 110, 22, 9; *texana* (Cret.-Texas), 2(4) : 85, 11, 8; *trinitatensis* (Cret.-Trinidad), 4(4) : 95, 13, 13.
- Truncatulina* cf. *basiloba* (Mioc.-Calif.), 2(3) : 62; *excolata* (Cret.-Mexico), 2(1) : 22, 3, 2; *spinea* (Cret.-Mexico), 2(1) : 22, 2, 10; *velascoensis* (Cret.-Mexico), 1(1) : 20, 3, 2.
- Tubulogenerina**, 2(4) : 78; *aperta* (Olig.-Miss.), 5(2) : 45, 8, 1.
- Turrispirillina**, 3(1) : 73.
- Turritellella grandis* (Penn.-Texas), 3(3) : 149, 26, 9; *spirans* (Penn.-Michigan), 3(2) : 109, 22, 5, 6.
- Uvigerina auberiana* (Tert.-Venezuela), 5(4) : 95, 13, 36; *beccarii* (Tert.-Venezuela), 5(4) : 95, 13, 37; *cocoaensis* (Eoc.-Ala.), 1(3) : 68, 10, 12; *compressa* (Mioc.-Austria), 1(1) : 10, 4, 2; *curta* (Eoc.-Trinidad), 5(1) : 13, 3, 13-15; *gallowayi* (Tert.-Ecuador), 5(4) : 94, 13, 33, 34; *hispidula* (Tert.-Venezuela), 5(4) : 95, 13, 35; *jacksonensis* (Eoc.-Ala.), 1(3) : 67, 10, 13; *seligi* (Cret.-La.), 1(1) : 1, 4, 1; *seriata* (Tert.-Trinidad), 5(1) : 13, 3, 11, 12; *spinicostata* (Tert.-Trinidad), 5(1) : 12, 3, 9, 10; *topilensis* (Eoc.-Mexico), 1(1) : 5, 1, 5.
- Uvigerina* (*Uvigerinella*) *californica* (Mioc.-Calif.), 2(3) : 58, 8, 2, 5; var. *ornata* (Mioc.-Calif.), 2(3) : 59, 8, 1, 6; *nudo-costata* (Mioc.-Calif.), 2(3) : 59, 8, 4, 8; *obesa* (Mioc.-Calif.), 2(3) : 59, 8, 3, 7.

- Uvigerinella**, 2(3) : 58; **parva** (Eoc.-Trinidad), 5(1) : 12, 3, 7.
- Vaginulina senticosa* (Eoc.-Trinidad), 5(1) : 7, 2, 5, 6.
- Valvulina ocalana* (Eoc.-Fla.), 2(2) : 34, 5, 4.
- Valvulineria**, 2(3) : 59; **californica** (Mioc.-Calif.), 2(3) : 60, 9, 1; var. **appressa** (Mioc.-Calif.), 2(3) : 60, 9, 5; var. **obesa** (Mioc.-Calif.), 2(3) : 61, 9, 2; **miocenica** (Mioc.-Calif.), 2(3) : 61, 8, 9, 10; 9, 3; var. **depressa** (Mioc.-Calif.), 2(3) : 61, 9, 7; **ornata** (Mioc.-Calif.), 2(3) : 61, 9, 4; *vilardeboana* (Mioc.-Calif.), 2(3) : 62, 9, 6.
- [Venezuela]. A Late Tertiary Fauna of Venezuela and Other Related Regions, 5(4) : 77.
- Ventilabrella**, 4(1) : 2; **eggeri** (Cret.), 4(1) : 2, 1, 10-12.
- Verneuilina sculptilis* (Eoc.-Ala.), 2(2) : 34, 5, 3.
- Vertebralina*, The Genera *Renulina* and, 3(3) : 158.
- Virgulina gunteri* Cushman—A Correction of Name, 5(4) : 105.
- Virgulina bramletti* (Tert.-Venezuela), 5(4) : 94, 13, 30; **californiensis** (Mioc.-Calif.), 1(2) : 32, 5, 11; 2(3) : 54; **floridana** (Mioc.-Fla.), 5(3) : 54, 9, 7-10; **gunteri** (Mioc.-Fla.), 5(4) : 105; **pertusa**, 5(3) : 53, 9, 1-4.
- Vitriwebbina*, 3(4) : 189; 4(1) : 20.
- Vulvulina advena* (Eoc.-Ala.), 2(2) : 32, 4, 9; *capreolus* (Tert.-Ecuador), 5(4) : 80, 12, 6; **spinosa** (Alazan-Mexico), 3(2) : 111, 23, 1.

